

**COLD AND FLU DAYTIME/MULTI-SYMPTOM/NON-DROWSY- acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled
Walgreen Company**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Walgreens 44-659-Delisted

***Active ingredients
(in each liquid-filled capsule)***

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - sore throat
 - nasal congestion
 - headache
 - fever
 - minor aches and pains
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- liver disease
- thyroid disease
- diabetes
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- heart disease
- high blood pressure
- cough that occurs with too much phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- do not take more than 8 capsules per 24 hours
- adults and children 12 years and over: take 2 capsules with water every 4 hours
- children under 12 years: ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from heat, humidity and light
- see end flap for expiration date and lot number

Inactive ingredients

edible white ink, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol

Questions or comments?

1-800-426-9391

Principal Display Panel**Walgreens**

Compare to Vicks® DayQuil® Cold & Flu
LiquiCaps® active ingredients††

NDC 0363-6590-21

DAYTIME • NON-DROWSY

Cold & Flu

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

MULTI-SYMPTOM

- Relieves aches, fever, sore throat,
nasal congestion & cough

16

LIQUID CAPS

ACTUAL SIZE

**TAMPER EVIDENT: DO NOT USE IF
PACKAGE IS OPENED OR IF BLISTER**

**UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING**

Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey

††This product is not manufactured or
distributed by The Procter & Gamble
Company, owner of the registered trademark
Vicks® DayQuil® Cold & Flu LiquiCaps®.
50844 REV0418B65921

**DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015**

***Walgreens* _____**
100% SATISFACTION
GUARANTEED

walgreens.com ©2017 Walgreen Co.

**PRODUCT OF CHINA. PACKAGED AND
QUALITY ASSURED IN THE U.S.A.**

Walgreens

DAYTIME • NON-DROWSY

Cold & Flu

Walgreens

DAYTIME • NON-DROWSY

Cold & Flu

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

MULTI-SYMPTOM

- Relieves aches, fever, sore throat, nasal congestion & cough

16

LIQUID CAPS



ACTUAL SIZE

Compare to Vicks® DayQuil® Cold & Flu LiquiCaps® active ingredients!

NDC 0363-6590-21

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Purpose

Acetaminophen 325 mg Pain reliever/fever reducer

Dextromethorphan HBr 10 mg Cough suppressant

Phenylephrine HCl 5 mg Nasal decongestant

Temporarily relieves common cold and flu symptoms:

■ sore throat ■ nasal congestion

■ headache ■ fever ■ minor aches and pains

■ cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

more than 4,000 mg of acetaminophen in 24 hours

■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: ■ skin redness ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2

days, is accompanied or followed by fever, headache, rash, nausea, or

vomiting, consult a doctor promptly.

Do not use

■ if you are now taking a prescription monoamine oxidase inhibitor

■ if you have ever had an allergic reaction to this product or any of its

ingredients

MAOI, ask a doctor or pharmacist before taking this product.

MAOI drug. If you do not know if your prescription drug contains an

conditions, or Parkinson's disease), or for 2 weeks after stopping the

■ if you have ever had an allergic reaction to this product or any of its

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Purpose

Acetaminophen 325 mg Pain reliever/fever reducer

Dextromethorphan HBr 10 mg Cough suppressant

Phenylephrine HCl 5 mg Nasal decongestant

Temporarily relieves common cold and flu symptoms:

■ sore throat ■ nasal congestion

■ headache ■ fever ■ minor aches and pains

■ cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

more than 4,000 mg of acetaminophen in 24 hours

■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: ■ skin redness ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2

days, is accompanied or followed by fever, headache, rash, nausea, or

vomiting, consult a doctor promptly.

Do not use

■ if you are now taking a prescription monoamine oxidase inhibitor

■ if you have ever had an allergic reaction to this product or any of its

ingredients

MAOI, ask a doctor or pharmacist before taking this product.

MAOI drug. If you do not know if your prescription drug contains an

conditions, or Parkinson's disease), or for 2 weeks after stopping the

■ if you have ever had an allergic reaction to this product or any of its

Drug Facts (continued)

Other information

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED

ON BLISTER IS TORN OR BROKEN

■ store at 25°C (77°F); excursions permitted between 15°-30°C

■ protect from heat, humidity and light

■ see end flap for expiration date and lot number

Drug Facts (continued)

Inactive ingredients

edible white ink, FD & C red #40, FD & C

yellow #6, gelatin, glycerin, polyethylene glycol, polyvinylpyrrolidone, propylene

glycol, purified water, sorbitol

Questions or comments? 1-800-426-9391

QUESTIONS OR COMMENTS? 1-800-426-9391

Directions

■ do not take more than directed

■ adults and children 12 years and over: take 2 capsules with water

■ children under 12 years: ask a doctor

every 4 hours

Keep out of reach of children. In case of accidental overdose, get

medical help or contact a Poison Control Center (1-800-222-1222) right

away. Prompt medical attention is critical for adults as well as for

children even if you do not notice any signs or symptoms.

■ pregnant or breast-feeding, ask a health professional before use.

could be signs of a serious condition.

■ fever gets worse or lasts more than 3 days

■ redness or swelling is present ■ new symptoms occur

■ cough comes back or occurs with rash or headache that lasts. These

■ Stop use and ask a doctor if

■ When using this product do not exceed recommended dosage.

■ Ask a doctor or pharmacist before use if you are taking the blood

thinning drug warfarin.

■ difficulty in urination due to enlargement of the prostate gland

■ cough that occurs with too much phlegm (mucus)

■ emphysema ■ heart disease

■ persistent or chronic cough such as occurs with smoking, asthma, or

■ liver disease ■ thyroid disease ■ diabetes

■ Ask a doctor before use if you have

acetaminophen, ask a doctor or pharmacist

■ with any other drug containing acetaminophen (prescription or

nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist

■ Stop use and ask a doctor if

■ When using this product do not exceed recommended dosage.

■ Ask a doctor or pharmacist before use if you are taking the blood

thinning drug warfarin.

■ difficulty in urination due to enlargement of the prostate gland

■ cough that occurs with too much phlegm (mucus)

■ emphysema ■ heart disease

■ persistent or chronic cough such as occurs with smoking, asthma, or

■ liver disease ■ thyroid disease ■ diabetes

■ Ask a doctor before use if you have

acetaminophen, ask a doctor or pharmacist

■ with any other drug containing acetaminophen (prescription or

nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist

■ Stop use and ask a doctor if

■ When using this product do not exceed recommended dosage.

■ Ask a doctor or pharmacist before use if you are taking the blood

thinning drug warfarin.

■ difficulty in urination due to enlargement of the prostate gland

■ cough that occurs with too much phlegm (mucus)

■ emphysema ■ heart disease

■ persistent or chronic cough such as occurs with smoking, asthma, or

■ liver disease ■ thyroid disease ■ diabetes

■ Ask a doctor before use if you have

acetaminophen, ask a doctor or pharmacist

■ with any other drug containing acetaminophen (prescription or

nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist

■ Stop use and ask a doctor if

■ When using this product do not exceed recommended dosage.

■ Ask a doctor or pharmacist before use if you are taking the blood

thinning drug warfarin.

■ difficulty in urination due to enlargement of the prostate gland

■ cough that occurs with too much phlegm (mucus)

■ emphysema ■ heart disease

■ persistent or chronic cough such as occurs with smoking, asthma, or

■ liver disease ■ thyroid disease ■ diabetes

■ Ask a doctor before use if you have

acetaminophen, ask a doctor or pharmacist

■ with any other drug containing acetaminophen (prescription or

nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist

■ Stop use and ask a doctor if

■ When using this product do not exceed recommended dosage.

■ Ask a doctor or pharmacist before use if you are taking the blood

thinning drug warfarin.

■ difficulty in urination due to enlargement of the prostate gland

■ cough that occurs with too much phlegm (mucus)

■ emphysema ■ heart disease

■ persistent or chronic cough such as occurs with smoking, asthma, or

■ liver disease ■ thyroid disease ■ diabetes

■ Ask a doctor before use if you have

acetaminophen, ask a doctor or pharmacist

■ with any other drug containing acetaminophen (prescription or

nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist

■ Stop use and ask a doctor if

■ When using this product do not exceed recommended dosage.

■ Ask a doctor or pharmacist before use if you are taking the blood

thinning drug warfarin.

■ difficulty in urination due to enlargement of the prostate gland

■ cough that occurs with too much phlegm (mucus)

■ emphysema ■ heart disease

■ persistent or chronic cough such as occurs with smoking, asthma, or

■ liver disease ■ thyroid disease ■ diabetes

■ Ask a doctor before use if you have

acetaminophen, ask a doctor or pharmacist

■ with any other drug containing acetaminophen (prescription or

nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist

■ Stop use and ask a doctor if

■ When using this product do not exceed recommended dosage.

■ Ask a doctor or pharmacist before use if you are taking the blood

thinning drug warfarin.

■ difficulty in urination due to enlargement of the prostate gland

■ cough that occurs with too much phlegm (mucus)

■ emphysema ■ heart disease

■ persistent or chronic cough such as occurs with smoking, asthma, or

■ liver disease ■ thyroid disease ■ diabetes

■ Ask a doctor before use if you have

acetaminophen, ask a doctor or pharmacist

■ with any other drug containing acetaminophen (prescription or

nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist

■ Stop use and ask a doctor if

■ When using this product do not exceed recommended dosage.

■ Ask a doctor or pharmacist before use if you are taking the blood

thinning drug warfarin.

■ difficulty in urination due to enlargement of the prostate gland

■ cough that occurs with too much phlegm (mucus)

■ emphysema ■ heart disease

■ persistent or chronic cough such as occurs with smoking, asthma, or

■ liver disease ■ thyroid disease ■ diabetes

■ Ask a doctor before use if you have

acetaminophen, ask a doctor or pharmacist

■ with any other drug containing acetaminophen (prescription or

nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist

■ Stop use and ask a doctor if

■ When using this product do not exceed recommended dosage.

■ Ask a doctor or pharmacist before use if you are taking the blood

thinning drug warfarin.

■ difficulty in urination due to enlargement of the prostate gland

■ cough that occurs with too much phlegm (mucus)

■ emphysema ■ heart disease

■ persistent or chronic cough such as occurs with smoking, asthma, or

■ liver disease ■ thyroid disease ■ diabetes

■ Ask a doctor before use if you have

acetaminophen, ask a doctor or pharmacist

■ with any other drug containing acetaminophen (prescription or

nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist

■ Stop use and ask a doctor if

■ When using this product do not exceed recommended dosage.

■ Ask a doctor or pharmacist before use if you are taking the blood

thinning drug warfarin.

■ difficulty in urination due to enlargement of the prostate gland

■ cough that occurs with too much phlegm (mucus)

■ emphysema ■ heart disease

■ persistent or chronic cough such as occurs with smoking, asthma, or

■ liver disease ■ thyroid disease ■ diabetes

■ Ask a doctor before use if you have

acetaminophen, ask a doctor or pharmacist

■ with any other drug containing acetaminophen (prescription or

nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist

■ Stop use and ask a doctor if

■ When using this product do not exceed recommended dosage.

■ Ask a doctor or pharmacist before use if you are taking the blood

thinning drug warfarin.

■ difficulty in urination due to enlargement of the prostate gland

■ cough that occurs with too much phlegm (mucus)

■ emphysema ■ heart disease

■ persistent or chronic cough such as occurs with smoking, asthma, or

■ liver disease ■ thyroid disease ■ diabetes

■ Ask a doctor before use if you have

acetaminophen, ask a doctor or pharmacist

■ with any other drug containing acetaminophen (prescription or

nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist

■ Stop use and ask a doctor if

■ When using this product do not exceed recommended dosage.

■ Ask a doctor or pharmacist before use if you are taking the blood

thinning drug warfarin.

■ difficulty in urination due to enlargement of the prostate gland

■ cough that occurs with too much phlegm (mucus)

■ emphysema ■ heart disease

■ persistent or chronic cough such as occurs with smoking, asthma, or

■ liver disease ■ thyroid disease ■ diabetes

■ Ask a doctor before use if you have

acetaminophen, ask a doctor or pharmacist

■ with any other drug containing acetaminophen (prescription or

nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist

■ Stop use and ask a doctor if

■ When using this product do not exceed recommended dosage.

■ Ask a doctor or pharmacist before use if you are taking the blood

thinning drug warfarin.

■ difficulty in urination due to enlargement of the prostate gland

■ cough that occurs with too much phlegm (mucus)

■ emphysema ■ heart disease

■ persistent or chronic cough such as occurs with smoking, asthma, or

■ liver disease ■ thyroid disease ■ diabetes

■ Ask a doctor before use if you have

acetaminophen, ask a doctor or pharmacist

■ with any other drug containing acetaminophen (prescription or

nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist

■ Stop use and ask a doctor if

■ When using this product do not exceed recommended dosage.

■ Ask a doctor or pharmacist before use if you are taking the blood

thinning drug warfarin.

■ difficulty in urination due to enlargement of the prostate gland

■ cough that occurs with too much phlegm (mucus)

■ emphysema ■ heart disease

■ persistent or chronic cough such as occurs with smoking, asthma, or

■ liver disease ■ thyroid disease ■ diabetes

■ Ask a doctor before use if you have

acetaminophen, ask a doctor or pharmacist

■ with any other drug containing acetaminophen (prescription or

nonprescription). If you are not sure whether a drug contains

acetaminophen, ask a doctor or pharmacist

■ Stop use and ask a doctor if

■ When using this product do not exceed recommended dosage.

COLD AND FLU DAYTIME/MULTI-SYMPTOM/NON-DROWSY

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-6590
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	

Product Characteristics

Color	orange (clear)	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	659
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-6590-08	2 in 1 CARTON	03/01/2015	09/09/2022
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0363-6590-22	4 in 1 CARTON	03/01/2015	05/11/2018

2		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:0363-6590-21	2 in 1 CARTON	03/01/2015	09/09/2022
3		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	03/01/2015	09/09/2022

Labeler - Walgreen Company (008965063)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	pack(0363-6590)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(0363-6590)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	pack(0363-6590)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(0363-6590)

Revised: 11/2021

Walgreen Company